

**β -AMYLOID PEPTIDE-BINDING PROTEINS
AND POLYNUCLEOTIDES ENCODING THE SAME**

5 This application claims benefit of U.S. Ser. No. 09/172,990 filed
October 14, 1998, U.S. Ser. No. 09/060, 609 filed April 15, 1998, now
abandoned, and U.S. Provisional Application 60/064,583, filed April 16,
1997, the content of which is incorporated into this application by
reference.

10

Field of the Invention

The present invention relates to a novel polynucleotides and
proteins encoded by such polynucleotides, along with therapeutic,
diagnostic, and research utilities for these polynucleotides and proteins.

15 In particular, the invention relates to polynucleotides and proteins encoded
by such polynucleotides which bind to β -amyloid peptide, one of the
primary components of amyloid deposits associated with Alzheimer's
Disease.

Background of the Invention

20 Alzheimer's disease (AD) is a progressive dementing disorder of
the elderly characterized by a series of structural abnormalities of the
brain. Neurons in multiple regions of the central nervous system (CNS)
become dysfunctional and die, resulting in alterations in synaptic inputs.
Cell bodies and proximal dendrites of these vulnerable neurons contain
25 neurofibrillary tangles composed of paired helical filaments, the main
component of which is a phosphorylated microtubular-binding protein,
namely tau. One of the hallmarks of the disease is the accumulation of
amyloid containing deposits within the brain called senile (or neuritic)
plaques. The principal component of amyloid plaques is β -amyloid peptide
30 (hereinafter "BAP", also referred in the literature as A β , β AP, etc.) which
forms dense aggregates during the course of AD.

BAP is a 39-43 amino acid peptide derived by proteolytic cleavage of amyloid precursor protein (hereinafter "APP") and composed of a portion of the transmembrane domain and the luminal/extracellular domain of APP. It is thought that the BAP peptide comprising 42 amino acids (BAP42) is potentially the more toxic aggregated form in humans. APP occurs as several BAP-containing isoforms. The major forms are comprised of 695, 751, and 770 amino acids, with the latter two APP containing a domain that shares structural and functional homologies with Kunitz serine protease inhibitors. In normal individuals, BAP does not accumulate and is rapidly removed from circulating fluids. However, the peptide can form plaques on surfaces of dystrophic dendrites and axons, microglia, and reactive astrocytes. The aggregation and deposition of BAP in neuritic plaques is postulated as one of the initiating events of AD. Investigation of the events leading to the expression and consequences of BAP and their individual roles in AD is a major focus of neuroscience research. In particular, the discovery of proteins that bind BAP is critical to advance understanding of the pathogenesis of the disease and to potentially introduce novel therapeutic targets.

Until the present invention, proteins and fragments thereof which bind with human BAP and which may be involved in the biological effects of BAP in AD had not been identified.

Summary of the Invention

This invention provides novel isolated polynucleotides which encode gene products that selectively bind human β -amyloid peptide (BAP) amino acid sequences.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) polynucleotide comprising the nucleotide sequence of SEQ ID NO: 1;
- (b) a polynucleotide comprising the nucleotide sequence of a β -

amyloid peptide-binding protein (BBP) of clone BBP1-fl deposited under
accession number ATCC 98617;

(c) a polynucleotide encoding a β -amyloid peptide-binding
protein (BBP) encoded by the cDNA insert of clone BBP1-fl deposited
5 under accession number ATCC 98617;

(d) a polynucleotide comprising the nucleotide sequence of SEQ
ID NO: 1 from nucleotide 202 to nucleotide 807;

(e) a polynucleotide comprising the nucleotide sequence of a β -
amyloid peptide-binding protein (BBP) of clone pEK196 deposited under
10 accession number ATCC 98399;

(f) a polynucleotide encoding a β -amyloid peptide-binding
protein (BBP) encoded by the cDNA insert of clone pEK196 deposited
under accession number ATCC 98399;

(g) a polynucleotide encoding a protein comprising the amino
15 acid sequence of SEQ ID NO: 2;

(h) a polynucleotide encoding a protein comprising a fragment
of the amino acid sequence of SEQ ID NO: 2 having human β -amyloid
peptide binding activity, the fragment comprising the amino acid sequence
from amino acid 68 to amino acid 269 of SEQ ID NO: 2;

(j) a polynucleotide which is an allelic variant of the
20 polynucleotide of (a)-(f) above;

(k) a polynucleotide which encodes a species homologue of the
protein of (g)-(i) above; and

(l) a polynucleotide capable of hybridizing under stringent
25 conditions to any one of the polynucleotides specified in (a)-(h).

Preferably such polynucleotide comprises the nucleotide sequence
of SEQ ID NO: 1; the nucleotide sequence of a β -amyloid peptide-binding
protein (BBP) of clone BBP1-fl deposited under accession number ATCC
98617; or a polynucleotide encoding a β -amyloid peptide-binding protein
30 (BBP) encoded by the cDNA insert of clone BBP1-fl deposited under

accession number ATCC 98617. Another embodiment provides the gene corresponding to the cDNA sequence of SEQ ID NO: 1.

- In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO: 2;
 - (b) the amino acid sequence of SEQ ID NO: 2 from amino acid 68 to amino acid 269;
 - (c) the amino acid sequence encoded by the cDNA insert of clone BBP1-fl deposited under accession number ATCC 98617; and
 - (d) fragments of the amino acid sequence of SEQ ID NO: 2 comprising the amino acid sequence from amino acid 185 to amino acid 217 of SEQ ID NO: 2.

Preferably such protein comprises the amino acid sequence of SEQ ID NO: 2 or the amino acid sequence of SEQ ID NO: 2 from amino acid 68 to amino acid 269. Fusion proteins are also claimed in the present invention.

In certain preferred embodiments, the polynucleotide is operably linked to an expression control sequence. The invention also provides a host cell, including bacterial, yeast, insect, and mammalian cells, transformed with such polynucleotides compositions.

Processes are also provided for producing a BBP which comprises (a) growing a culture of the host cell of claim 3 in a suitable culture medium; and (b) purifying the protein from the culture medium.

Compositions comprising an antibody which specifically reacts with such BBPs are also provided by the present invention.

Methods and diagnostic processes are provided for detecting a disease state characterized by the aberrant expression of human BAP, as well as methods for identifying compounds which regulate the activity of BBPs.

Another embodiment of the invention includes transgenic animals comprising a polynucleotide encoding a BBP operably linked to an expression control sequence.

Brief Description of the Drawings

5 The following drawings depict certain embodiments of the invention. They are illustrative only and do not limit the invention otherwise disclosed herein.

Figure 1: Yeast 2-hybrid screen design. A Y2H host strain expressing the Gal4 DNA-binding domain fused to BAP₄₂ (BAP^{BD}; plasmid containing TRP1 marker) and nonfusion BAP₄₂ (BAP; plasmid containing URA3 marker) was transformed with a Y2H human fetal brain cDNA library (plasmid containing LEU2 marker) expressing Gal4 activation domain fusion proteins (unknown^{AD}) as described. Therefore, strains
10 contained three episomal plasmids, denoted by circles, expressing the indicated protein. Positive protein-protein interactions reconstituted Gal4 activity at the upstream activating sequence (GALUAS) thereby inducing transcription of the reporter gene HIS3.

Figure 2: Demonstration of BBP1/BAP association. Y2H strains were assayed for histidine prototrophy by making 10-fold serial dilutions and spotting 5 μ l on synthetic agar medium lacking tryptophan, leucine,
20 histidine and containing 25 mM 3-amino-triazole as described. All strains contain the BAP fusion protein expression plasmid pEK162 as indicated by the label BAP. The first columns (vector) contain independently derived strains carrying pEK162 and the vector pACT2 expressing an irrelevant fusion protein. These serve as a measure of background for
25 comparison with strains expressing target proteins. The columns marked by BBP1Dtm express a truncated BBP1 from pEK198, as described in the text. The interaction between BAP and BBP1 Δ tm fusion proteins reconstitutes Gal4 activity, resulting in induction of a HIS3 reporter gene
30 (see Figure 1), observed as enhanced prototrophic growth compared to the control strains.

Figure 3: Bioassays demonstrating BBP1 interactions with G α proteins. The predicted intracellular domain of BBP1 was expressed as a Gal4 DNA-binding domain with portions of rat G α s, G α o, or G α i2 expressed as Gal4 activation domain fusion proteins. Y2H responses of two independently derived clones of each strain were compared to responses of cells lacking a G protein component (vector). The protocol is as described in the legend to Figure 2.

Figure 4: Localization of the interactions between BBP1 and BAP. BBP1 Δ tm was divided into two overlapping segments as described in the text. These proteins, BBP1 Δ C or BBP1 Δ N, were assayed for interactions with BAP. The assay method and the strains labeled vector or BBP1 Δ tm are as described in the legend to Figure 2. Strains labeled BBP1 Δ C or BBP1 Δ N express the indicated BBP1 segment as a fusion protein.

Figure 5: Expression of BBP1 mRNA in human tissues (A) and brain regions (B). Nylon membranes blotted with 2 μ g size fractionated poly-A RNA isolated from the indicated tissues were obtained from CLONTECH. These were hybridized with a radiolabeled BBP1 cDNA probe as described. A predominant band corresponding to 1.25 kb (determined from molecular weight markers, not shown) was observed in all lanes. Higher molecular weight bands likely correspond to heteronuclear RNA; the BBP1 gene contains several introns. Blots were stripped and reprobed with β -actin as a loading and RNA integrity control; all lanes exhibited equivalent signal (data not shown).

Figure 6: Expression of BBP1 and APP in cells of the hippocampus. Images of in situ hybridization autoradiograms showing the pattern of BBP1 (A) and APP (B) expression in human hippocampal and entorhinal cortex. The sections used to generate these images were taken from postmortem specimens obtained from two different patients. Abbreviations: DG = dentate gyrus; CA1 = hippocampal subfield; EC = entorhinal cortex.

Figure 7: Comparison of BBP1 interactions with human or rodent BAP. Rodent BAP was engineered and expressed as a fusion protein as described in the text. The strains labeled human BAP are identical to those shown in Figure 2. The strains labeled rodent BAP express rodent BAP as the Gal4 DNA-binding domain fusion. Vector indicates control strains containing only vector opposing the BAP fusion proteins; BBP1 indicates strains expressing the BBP1 Δ tm fusion protein.

Figure 8. In vitro BBP1 binding to beta-amyloid protein (1-42). In vitro transcribed and translated N-terminally myc-tagged BBP1 (~20ng) was incubated with disaggregated human [125 I]-Tyr-BAP₍₁₋₄₂₎. The myc-tagged BBP1 was immunoprecipitated with mouse anti-myc antibody and rabbit anti-mouse IgG antibody conjugated to protein A-Agarose beads. lane 1, MW markers; lane2, 4, 6, 8: blank; lane 3: [125 I]-Tyr-BAP₍₁₋₄₂₎(disaggregated; \approx 25nCi); lane 5: Pre-incubation of rabbit anti-mouse IgG antibody conjugated to protein A-Agarose beads in Superblock, plus mouse anti-myc antibody and radiolabeled BAP (no BBP1); lane 7: Pre-incubation of rabbit anti-mouse IgG antibody conjugated to protein A-Agarose beads in low salt binding buffer (see Material and Methods), plus mouse anti-myc antibody and radiolabeled BAP (no BBP1); lane9: Pre-incubation of rabbit anti-mouse IgG antibody conjugated to protein A-Agarose beads in Superblock, plus mouse anti-myc antibody, radiolabeled BAP, and in vitro synthesized myc-tagged BBP1.

Figure 9. Effect of BBP1 on Nt2 Stem Cells Exposed to BAP. Nt2 neurons, and Nt2 stem cells transfected with pEGFP alone or cotransfected with pEGFP plus pBBP1 (also referred to as pOZ363), were treated for 48 hrs with aged (toxic) or fresh (nontoxic) BAP preparations as described in the text. Final concentrations were 2 μ M or 5 μ M, respectively. Cells were prepared for determinations of nuclear morphology as described. Values represent the mean percent condensed nuclei, derived from viewing multiple fields for each sample. For the stem cell populations, only transfected cells (EGFP+) were scored.

Figure 10. Sensitivity of Nt2 stem cells to modified BBP. Nt2 stem cells were transfected with pEGFP alone or cotransfected with pEGFP plus BBP1 or BBP1(R138E) expression plasmids. Samples were treated with BAP at the indicated concentrations for 44 hrs. Cells were prepared for determinations of nuclear morphology as described. Values represent percent-condensed nuclei of transfected cells (EGFP+). The differences between values for BBP1 samples versus controls at 2 or 8 μ M BAP are highly significant ($P < 0.005$) as determined by a Yates modified chi-square test of probability.

Detailed Description of the invention

The present invention relates to the isolation and cloning of a human β -amyloid peptide binding protein (BBP1). BBP1 has been characterized as a fusion protein in a yeast 2 hybrid assay as binding to BAP, specifically the 42 amino acid fragment of BAP (BAP42). Expression of BBP1 has been shown in human tissues and in specific brain regions (Figure 5). Importantly BBP1 has been demonstrated to selectively bind human BAP in a yeast 2 hybrid system as compared to rodent BAP. These findings support the premise that the BBP1 of the present invention may be used in the diagnosis and treatment of Alzheimer's Disease.

The BBP1 Coding Sequence

The initial human BBP1 clone (designated clone 14) was obtained by using a yeast 2-hybrid (Y2H) genetic screen developed to identify proteins which interact with human BAP₄₂, a potentially more toxic form of BAP. BAP₄₂ was expressed fused to the yeast Gal4 DNA-binding domain and was also expressed as free peptide (Figure 1). This strain was transformed with a human fetal brain cDNA Y2H library. A single clone, denoted #14, from approximately 10^6 independent transformants, produced consistent reporter gene activation and contained a substantial open reading frame continuous with that of the GAL4 domain. The cDNA

insert comprised 984 base pairs, terminating in a poly-A tract. This sequence encoded 201 amino acids (amino acid 68 to amino acid 269 of SEQ ID NO: 2) with two regions of sufficient length and hydrophobicity to transverse a cellular membrane. There are also potential asparagine-linked glycosylation sites. Clone 14 was designated clone pEK196 and was deposited with American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110-2209, on April 9, 1997 and assigned Accession Number 98399. All deposits referred to in this application refer to deposits with ATCC and all such deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and under conditions that will make them available to the public as of the issue date of any patent granted from this application.

The library-derived plasmid was isolated from clone 14 and used to reconstruct Y2H assay strains. Examination of these strains demonstrated that the BAP fusion protein specifically interacted with the clone 14 protein, although the response was weak. Since protein domains of strong hydrophobicity, such as transmembrane regions, inhibit Y2H responses (Ozenberger, unpublished data), clone 14 insert was truncated (BBP1 Δ tm; see Table 2 below for further description) to remove the region of strongest hydrophobicity and retested for interactions with BAP. A much more robust Y2H response was observed with BBP1 Δ tm, supporting the notion that the deleted sequences encode a potential transmembrane ("tm") anchor. Clone 14 identifies a novel BAP binding protein in the form of a fusion protein.

The BBP1 cDNA sequences contained in clone 14 were identified as lacking the 5' end of the protein coding region as no potential initiating methionine codon was present. Multiple attempts at conventional 5' RACE (rapid amplification of cDNA ends) utilizing a standard reverse-transcriptase only resulted in the addition of 27 nucleotides. Thus, a

genomic cloning approach as described in Example 2, below, was used to isolate the 5' terminus.

Since the 5' coding sequence terminus was derived from a genomic library, there existed the possibility that this region contained
5 introns. This potentiality was investigated by two methods as described in Example 2, below. The resulting data confirmed the upstream sequences (both from genomic and cDNA sources) and the lack of introns in this region. Plasmid BBP1-fl containing a cDNA insert encoding the full length BBP1 protein coding region was deposited in the American Type
10 Culture Collection with accession number 98617 on December 11, 1997. The entire coding region and deduced protein sequence is shown in SEQ ID NOS:1 and 2..The 3' nontranslated nucleotide sequences are contained in the original clone 14 (pEK196).

In accordance with the present invention, nucleotide sequences
15 which encode BBP1, fragments, fusion proteins or functional equivalents thereof, may be used to generate recombinant DNA molecules that direct the expression of BBP1, or a functionally active peptide, in appropriate host cells. Alternatively, nucleotide sequences which hybridize to portions of the BBP1 sequence may be used in nucleic acid hybridization
20 assays, Southern and Northern blot assays, etc.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The present invention also includes polynucleotides capable of hybridizing under reduced stringency conditions, more preferably stringent
25 conditions, and most preferably highly stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced
30 stringency conditions are at least as stringent as, for example, conditions M-R.

Stringency Conditions

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) ¹	Hybridization Temperature and BufferH	Wash Temperature and BufferH
A	DNA:DNA	> 50	65EC; 1xSSC -or- 42EC; 1xSSC, 50% formamide	65EC; 0.3xSSC
B	DNA:DNA	< 50	T _B *; 1xSSC	T _B *; 1xSSC
C	DNA:RNA	> 50	67EC; 1xSSC -or- 45EC; 1xSSC, 50% formamide	67EC; 0.3xSSC
D	DNA:RNA	< 50	T _D *; 1xSSC	T _D *; 1xSSC

E	RNA:RNA	> 50	70EC; 1xSSC -or- 50EC; 1xSSC, 50% formamide	70EC; 0.3xSSC
F	RNA:RNA	< 50	T _F *; 1xSSC	T _F *; 1xSSC
G	DNA:DNA	> 50	65EC; 4xSSC -or- 42EC; 4xSSC, 50% formamide	65EC; 1xSSC
H	DNA:DNA	< 50	T _H *; 4xSSC	T _H *; 4xSSC
I	DNA:RNA	> 50	67EC; 4xSSC -or- 45EC; 4xSSC, 50% formamide	67EC; 1xSSC
J	DNA:RNA	< 50	T _J *; 4xSSC	T _J *; 4xSSC
K	RNA:RNA	> 50	70EC; 4xSSC -or- 50EC; 4xSSC, 50% formamide	67EC; 1xSSC
L	RNA:RNA	< 50	T _L *; 2xSSC	T _L *; 2xSSC
M	DNA:DNA	> 50	50EC; 4xSSC -or- 40EC; 6xSSC, 50% formamide	50EC; 2xSSC
N	DNA:DNA	< 50	T _N *; 6xSSC	T _N *; 6xSSC
O	DNA:RNA	> 50	55EC; 4xSSC -or- 42EC; 6xSSC, 50%	55EC; 2xSSC

			formamide	
P	DNA:RNA	< 50	T _P *; 6xSSC	T _P *; 6xSSC
Q	RNA:RNA	> 50	60EC; 4xSSC -or- 45EC; 6xSSC, 50% formamide	60EC; 2xSSC
R	RNA:RNA	< 50	T _R *; 4xSSC	T _R *; 4xSSC

l: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

^H: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

*T_B - T_R: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10EC less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, T_m(EC) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T_m(EC) = 81.5 + 16.6(log₁₀[Na⁺]) + 0.41(%G + C) - (600/N), where N is the number of bases in the hybrid, and [Na⁺] is the concentration of sodium ions in the hybridization buffer ([Na⁺] for 1xSSC = 0.165 M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and Current

Protocols in Molecular Biology, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that
5 is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it
10 hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

Expression of BBP1

15 The isolated polynucleotide of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., Nucleic Acids Res. 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of
20 expressing recombinant proteins are also known and are exemplified in R. Kaufman, Methods in Enzymology 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which
25 has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

Expression Systems for BBP1

A number of types of cells may act as suitable host cells for
30 expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney

293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

- 5 Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially
- 10 suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to
- 15 obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

- The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression
- 20 system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac7 kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by
- 25 reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

- The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be
- 30 purified from such culture (i.e., from culture medium or cell extracts) using known purification processes, such as gel filtration and ion

exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl7 or Cibacrom blue 3GA

- 5 Sepharose7; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

- Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed
10 as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLab (Beverly, MA), Pharmacia (Piscataway, NJ) and InVitrogen, respectively. The protein can also be tagged with an epitope
15 and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available from Kodak (New Haven, CT).

- Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media,
20 e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and
25 is defined in accordance with the present invention as an "isolated protein."

- The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells
30 containing a nucleotide sequence encoding the protein.

The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are known to those skilled in the art. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., USP No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

Yeast 2 Hybrid Assays

Y2H assays demonstrated that the association of BAP with the BBP1 fusion protein is specific. The association of BBP1 with BAP suggests that BBP1 activity may have a defined role in the pathogenesis of Alzheimer's disease.

BBP1 sequences were compared to Genbank using the basic local alignment search tool (BLAST; Altschul et al., 1990). The BBP1 protein and translations of available expressed sequence tags were aligned, searched for conserved segments, and evaluated by the MoST (Tatusov et al., 1994) protein motif search algorithm. These analyses revealed a potential evolutionary relationship to the G protein-coupled receptor (GPCR) family. Specifically, these analyses indicated that BBP1 contains two potential transmembrane (tm) domains equivalent to tm domains 3 and 4 of G protein-coupled receptors. The intervening hydrophilic loop contains a well-characterized three amino acid motif, aspartate (D) or glutamate followed by arginine (R) and an aromatic residue (Y or F) (commonly referred to as the DRY sequence), that is conserved in almost all members of this receptor family and has been shown to serve as a molecular trigger for G protein activation (Acharya and Karnik, 1996).

Data from Y2H assays (see Figures 2-4) indicate that BBP1 represents a novel protein potentially containing a functional module shared with members of the G protein-coupled receptor superfamily. Specifically, it appears that BBP1 retains the critical DRF sequence (amino acids 199 to amino acids 201 of SEQ ID NO: 2), between two predicted tm domains, and may have the potential to couple to a G protein regulated signaling pathway.

APP has been shown to functionally associate with Gao (Nishimoto et al., 1993; Yamatsuji et al., 1996) and BBP1 contains a structural motif known to be a G α protein activating sequence in the related G protein-coupled receptors. Additionally, a hypothesis based on the predicted position and orientation of BBP1 tm domains suggests that

the region of the protein that interacts with BAP would be topographically constrained to the same location as BAP in APP.

Y2H assay strains were engineered to evaluate the association of the BBP1 intracellular region with G α proteins. The predicted intracellular sequences of BBP1 were expressed as a fusion protein and assayed for interactions with C-terminal regions of three G α proteins. Protein segments used in these experiments are listed in Table 2, below. The BBP1 intracellular loop interacted with all three G α proteins (Figure 3), supporting the premise that BBP1 may function as a modulator of G protein activity. These various Y2H assays suggest the intriguing model of a multiple protein complex minimally composed of the integral membrane proteins BBP1 and APP coupled to a heterotrimeric G protein.

Table 2. Plasmids used in yeast 2-hybrid assays

expression plasmid	protein	segment
	BAP	
pEK162	(human)	1 - 42
pEK240	(mouse)	1 - 42
	BBP1	
pEK196	(clone 14)	68 - 269
pEK198	(Δ tm)	68 - 202
pEK219	(Δ C)	68 - 175
pEK216	(Δ N)	123 - 202
pOZ339	(intracellular)	185 - 217
	G α	
pOZ345	(G α s)	235 - 394
pOZ346	(G α o)	161 - 302
pOZ348	(G α i2)	213 - 355

Further analysis of BBP1 was obtained using Y2H assays. Two overlapping portions of the BBP1 sequences contained in the BBP1 Δ tm clone were amplified and cloned into the Y2H vector pACT2 (expression

plasmids pEK216 and pEK219, Table 2 and corresponding proteins BBP1 Δ N and BBP1 Δ C, (Figure 4)). The Δ C construct lacked both tm domains; the Δ N construct encoded the first tm domain plus the proceeding 52 amino acids. These fusion proteins were assayed with the BAP fusion protein and responses compared to those of strains expressing the larger BBP1 Δ tm protein. The BBP1 Δ C protein induced a weak Y2H response (compare BBP1 Δ C to vector, Figure 4), but the BBP1 Δ N protein, containing the first tm domain and adjacent amino-proximal sequences produced a response only slightly weaker than that observed with BBP1 Δ tm (Figure 4). These results suggest that a major determinant for the association with BAP is contained within the BBP1 region predicted to be topographically similar to BAP in the wild-type APP protein.

The Y2H system was utilized to demonstrate the selectivity and specificity of BBP1 binding to human BAP as compared to rodent BAP. There are three amino acid substitutions (G5R, F10Y and R13H) in the rodent BAP sequence compared to the human sequence. It was of interest, to evaluate the association of rodent BAP with BBP1 in the Y2H system. The sequence of human BAP in pEK162 was changed to encode the rodent peptide by oligonucleotide directed mutagenesis by PCR. The resultant plasmid, pEK240, is identical to the human BAP fusion protein expression plasmid utilized throughout this report except for the three codons producing the amino acid substitutions for the rodent peptide sequence. Interactions between BBP1 fusion protein and rodent and human BAP fusion proteins were compared by Y2H bioassay. Strains expressing BBP1 and the rodent BAP failed to produce a growth response (Figure 7). This finding supports the premise that BBP1 may serve as a specific mediator of the neurotoxic effects of BAP, and provides a mechanism to explain the reduced neurotoxicity of the rodent BAP. Importantly, these data also serve to illustrate the high degree of specificity of the BBP1/BAP interaction in the Y2H assays since the

substitution of three amino acids was sufficient to completely abrogate the association (Figure 7).

Isolated BBP1 polypeptides

5 Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95%
10 identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more
15 preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

Species homologues of the disclosed polynucleotides and proteins
20 are also provided by the present invention. As used herein, a species homologue is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide. Preferably, polynucleotide species homologues have at least 60% sequence identity
25 (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, and protein species homologues have at least 30% sequence identity (more preferably, at least 45% identity; most preferably at least 60% identity) with the given protein, where sequence identity is determined by comparing the nucleotide
30 sequences of the polynucleotides or the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while

minimizing sequence gaps. Species homologues may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species. Preferably, species homologues are those isolated from mammalian species. Most preferably, species homologues are those isolated from certain mammalian species such as, for example, *Pan troglodytes*, *Gorilla gorilla*, *Pongo pygmaeus*, *Hylobates concolor*, *Macaca mulatta*, *Papio papio*, *Papio hamadryas*, *Cercopithecus aethiops*, *Cebus capucinus*, *Aotus trivirgatus*, *Sanguinus oedipus*, *Microcebus murinus*, *Mus musculus*, *Rattus norvegicus*, *Cricetulus griseus*, *Felis catus*, *Mustela vison*, *Canis familiaris*, *Oryctolagus cuniculus*, *Bos taurus*, *Ovis aries*, *Sus scrofa*, and *Equus caballus*, for which genetic maps have been created allowing the identification of syntenic relationships between the genomic organization of genes in one species and the genomic organization of the related genes in another species (O'Brien and Seuanez, 1988, *Ann. Rev. Genet.* 22: 323-351; O'Brien et al., 1993, *Nature Genetics* 3:103-112; Johansson et al., 1995, *Genomics* 25: 682-690; Lyons et al., 1997, *Nature Genetics* 15: 47-56; O'Brien et al., 1997, *Trends in Genetics* 13(10): 393-399; Carver and Stubbs, 1997, *Genome Research* 7:1123-1137; all of which are incorporated by reference herein).

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotides which also encode proteins which are identical or have significantly similar sequences to those encoded by the disclosed polynucleotides. Preferably, allelic variants have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps. Allelic variants may be isolated and identified by making suitable probes or primers from the sequences provided herein

and screening a suitable nucleic acid source from individuals of the appropriate species.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

5

Applications

BBP1 proteins of the present invention can be used in a variety of applications routine to one of skill in the art based upon this disclosure. Specifically the BBPs can be used as immunogens to raise antibodies

10 which are specific to the cloned polypeptides. Various procedures known in the art may be used for the production of antibodies to BBP1 proteins. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments and a Fab expression library. For the production of antibodies, various host animals including, but not

15 limited to rabbits, mice, and rats, are injected with a BBP. In one embodiment, the polypeptide or a fragment of the polypeptide capable of specific immunoactivity is conjugated to an immunogenic carrier. Adjuvants may also be administered in conjunction with the polypeptide to increase the immunologic response of the host animal. Examples of

20 adjuvants which may be used include, but are not limited to, complete and incomplete Freund's, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol.

25 Monoclonal antibodies to BBP1 proteins of the present invention can be prepared using any technique which provides for the production of antibodies by continuous cell line in culture. Such techniques are well known to those of skill in the art and include, but are not limited to, the hybridoma technology originally described by Kohler and Milstein (Nature

30 1975, 256,4202-497), the human B-cell hybridoma technique described by Kosbor et al. (Immunology Today 1983, 4, 72) and the EBV-

hybridoma technique described by Cole et al. (Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp 77-96).

Antibodies immunoreactive to the polypeptides of the present invention can then be used to screen for the presence and subcellular
5 distribution of similar polypeptides in biological samples. In addition, monoclonal antibodies specific to the BBP1 proteins of the present invention can be used as therapeutics.

The BBP1 proteins can also serve as antigens useful in solid phase assays measuring the presence of antibodies which immunoreact with the
10 claimed peptides. Solid phase competition assays can be used to measure immunological quantities of clone 14-related antigen in biological samples. This determination is not only useful in facilitating the complete characterization of the cellular function or functions of the polypeptides of the present inventions, but can also be used to identify patients with
15 abnormal amounts of these proteins.

BBP1 proteins of the present invention can also be used as capture reagents in affinity chromatography for the detection of BAP and BAP aggregates as markers for AD.

In addition, these BBP1s are useful as reagents in an assay to
20 identify candidate molecules which effect the interaction of BAP and the cloned protein. Compounds that specifically block this association could be useful in the treatment or prevention of AD.

These BBP1s are also useful in acellular in vitro binding assays wherein alteration by a compound in the binding of these beta amyloid
25 peptide associated proteins to BAP or BAP aggregates is determined. Acellular assays are extremely useful in screening sizable numbers of compounds since these assays are cost effective and easier to perform than assays employing living cells. Upon disclosure of the polypeptides of the present invention, the development of these assays would be routine
30 to the skilled artisan. In such assays, either BBP1 or BAP is labeled. Such labels include, but are not limited to, radiolabels, antibodies, and

fluorescent or ultraviolet tags. Binding of a BBP1 to BAP or BAP aggregates is first determined in the absence of any test compound. Compounds to be tested are then added to the assay to determine whether such compounds alter this interaction.

5

Examples

The present invention is further described by the following examples. The examples are provided solely to illustrate the invention by reference to specific embodiments. These exemplifications, while
10 illustrating certain specific aspects of the invention do not portray the limitations or circumscribe the scope of the invention.

Yeast two-hybrid system (hereinafter "Y2H"): Y2H expression plasmids were constructed in vectors pAS2 and pACT2 (described in WadeHarper et al., 1993) and pCUP (described in Ozenberger and Young,
15 1995). Yeast strain CY770 (Ozenberger and Young, 1995) served as the host for all Y2H assays.

Genetic screen: The polymerase chain reaction (PCR) method was used to amplify and modify sequences encoding BAP. Oligonucleotides #1 (5' - CC ATG GAT GCA GAA TTC CGA C) and #3 (5' -
20 AAGCTTGTGCGAC TTA CGC TATGAC AAC ACC GC) were used to amplify BAP using pCLL621, a modified human APP clone (Jacobsen et al., 1994), as template. The amplified DNA consists of codons 389 to 430 (which encodes BAP₄₂) of the APP precursor protein with the following modifications. The sense strand primer added a 5' NcoI
25 restriction site in the same translational reading frame as the NcoI site in pAS2. The antisense strand primer added a stop codon and HindIII and Sall sites for cloning. The product from this amplification was ligated into the TA cloning system (Invitrogen Corp., Carlsbad, CA), and subsequently removed by digestion with NcoI and Sall. This fragment was cloned into
30 pAS2 cleaved with NcoI plus Sall. The resultant plasmid, pEK162, was confirmed by DNA sequencing through the GAL4/BAP junction. The

protein (BAP^{BD}; Figure 1) expressed from pEK162 comprised a fusion protein containing the DNA-binding domain of the yeast transcriptional activation protein Gal4 (lacking functional activation sequences) with the addition of the 42 amino acids of BAP to the carboxy-terminus. An
5 expression plasmid was developed that mediates the expression of unmodified BAP₄₂. Oligo #2 (5' - AAGCTTAAG ATG GAT GCA GAA TTC CGA C) was paired with oligo #3 in a PCR as described above. The product of this amplification contains a 5' HindIII site and translation initiation signals optimized for expression in *Saccharomyces cerevisiae*.
10 Again, the DNA fragment was cloned into the TA system. It was then isolated on a HindIII fragment and cloned into pCUP cleaved with HindIII. The orientation of the BAP gene in the resultant plasmid, pEK149 (BAP; Figure 1), was confirmed by DNA sequencing. The BAP expression plasmids pEK149 (which used URA3 as the selection marker) and pEK162
15 (which used TRP1 as the selection marker) were transformed into the yeast host CY770 (Ozenberger and Young, 1995). The strain containing both plasmids was designated CY2091. A plasmid library consisting of cDNA fragments isolated from human fetal brain cloned into the yeast 2-hybrid expression vector pACT2 (which used LEU2 as the selection
20 marker) was purchased from Clontech Laboratories, Inc. (Palo Alto, CA). The library-derived protein is depicted in Figure1 as unknown^{AD}. This library was used to transform CY2091. The samples were spread on synthetic complete (SC) yeast growth medium lacking uracil, tryptophan, and leucine to select cells containing all three plasmids. The medium also
25 lacked histidine and contained 3-amino-triazole, an inhibitor of the product of the yeast HIS3 gene, at a concentration of 25 mM. 3-Amino-triazole was utilized to reduce activity from low-level constitutive expression of the HIS3 reporter gene. Plates were incubated at 30°C for 12 days. Twenty-four colonies exhibiting increased histidine prototrophy were
30 isolated. Transformation controls indicated that the screen assayed 10⁶ individual clones. A PCR approach was utilized to quickly determine the

content of positive clones. Total DNA was isolated from each positive strain by standard methods. This material was used as template for PCRs using oligos #4 (5' - TTTAATACCA CTACAATGGA T) plus #5 (5' - TTTTCAGTAT CTACGATTCA T) which flank the cloning region of the library vector pACT2. DNA fragments were ligated into the TA system and examined by DNA sequencing. The library plasmid contained in clone #14 (as described above) was isolated by shuttle into *E. coli*. The nucleotide sequence of the human cDNA sequences was determined, confirming the sequence of the initial PCR product.

Bioassays: Strains were grown overnight in 2ml SC medium lacking leucine and tryptophan to a density of approximately 7×10^7 cells per ml. Cells were counted and 10-fold serial dilutions made from 10^4 to 10^8 cells per ml in sterile water. These samples were spotted in 5 μ l aliquots on SC medium lacking leucine, tryptophan and histidine and containing 25 mM 3-amino-triazole. Plates were incubated at 30°C for 2 to 3 days. Positive protein/protein interactions were identified by increased prototrophic growth compared to control strains expressing the Gal4 DNA-binding domain fusion protein plus an irrelevant transcriptional activation domain fusion protein (or simply containing the pACT vector without inserted sequences). These control strains were indicated in the Figures described above as the label 'vector'. This assay method was highly reproducible and provided for the detection of subtle inductions of growth mediated by the specific interaction between target proteins. The original BBP1 clone, designated pEK196 and deposited as ATCC 98399; is referred herein as clone 14), was used as a PCR template to truncate the protein product to express BBP1 Δ tm. Sense primer #6 (5'-TTTAATACCA CTACAATGGA T) annealed to GAL4 sequences in pACT2. The antisense primer #7 (5'-CTCGAG TTA AAA TCG ATC TGC TCC CAA CC) incorporated a 3' stop codon and XhoI site immediately 3' to the sequences encoding the DRF motif of BBP1. The PCR product was ligated into the TA cloning vector and subsequently digested with EcoRI

+ XhoI and cloned into pACT2. The hybrid product expressed from this plasmid (pEK198) was denoted BBP1 Δ tm. Similarly, primer #7 was paired with primer #8 (5'-GAATT CCA AAA ATA AAT GAC GCT ACG) to engineer the BBP1 Δ N expression plasmid pEK216. Again, the PCR

5 product was ligated into the TA system and the resultant plasmid digested with EcoRI + XhoI with the BBP1 fragment (codons 123-202) finally ligated into pACT2 digested with the same enzymes. BBP1 Δ C was made by using the pACT2-specific oligo #6 with antisense oligo #9 (5'-CTCGAG TCA AGA TAT GGG CTT GAA AAA AC). After TA cloning,

10 isolation of the EcoRI-XhoI fragment and cloning into pACT2, the resultant plasmid, pEK219, expressed BBP1 from residue 68 to 175. Sequences encoding the BBP1 intracellular loop were amplified using oligonucleotides #10 (5'-CCTTCC ATG GAA GTG GCA GTC GCA TTG TCT) plus #11 (5'-AACACTCGAG TCA AAA CCC TAC AGT GCA AAA C). This product,

15 containing BBP1 codons 185 to 217, was digested with NcoI + XhoI and cloned into pAS2 cleaved with NcoI + SalI to generate pOZ339. Construction of all G α protein expression plasmids utilized the BamHI site near the center of each rat cDNA sequence (Kang et al., 1990) as the site of fusion in pACT2. Sense primers annealed to sequences 5' of the

20 BamHI site; antisense primers annealed to sequences 3' of the stop codon and included a SalI restriction site. Primers were: G α_o , sense (#17) = 5'-GTGGATCCAC TGCTTCGAGG AT, antisense (#18) = 5'-GTCGACGGTT GCTATACAGG ACAAGAGG; G α_s , sense (#19) = 5'-GTGGATCCAG TGCTTCAATG AT, antisense (#20) = 5'-GTCGACTAAA

25 TTTGGGCGTT CCCTTCTT; G α_i2 , sense (#21) = 5'-GTGGATCCAC TGCTTTGAGG GT, antisense (#22) = 5'-GTCGACGGTC TTCTTGCCCC CATCTTCC. PCR products were cloned into the TA vector. G α sequences were isolated as BamHI-SalI fragments and cloned into pACT2 digested with BamHI + SalI. See Table 2 for plasmid designations.

30 Finally, oligonucleotide #23 was synthesized for the conversion of human BAP to the rodent sequence. This primer has the sequence 5'-

ATATGGCCATG GAT GCA GAA TTC GGA CAT GAC TCA GGA T
GAA GTT CGT. The triplets represent the first 13 codons of BAP; the
three nucleotides that were changed to produce the rodent sequence are
underlined. Oligo #23 was paired with #24 (5'-TGACCTACAG

- 5 GAAAGAGTTA) which anneals to a region of the Y2H vectors that is 3'
of the cloning site in a PCR using pEK162 as the template. The product
was cleaved with NcoI + Sall and ligated into pAS2 to produce pEK240.
The nucleotide sequence of the segment encoding rodent BAP was
confirmed.
- 10 Genomic cloning; RACE (rapid amplification of cDNA ends): A human
genomic lambda library (Stratagene), corresponding to $\approx 2.0 \times 10^6$ pfus,
was screened with randomly-primed EcoRI/ClaI fragment probe
corresponding to nucleotides 187-600 (Figure 2). The probe was labeled
with [32 P]-CTP using the 32 P QuickPrimer Kit according to the
- 15 manufacturer's (Pharmacia) protocol. Filters were hybridized under high
stringency: 40°C in 50% formamide, 0.12M NaHPO₄, 0.25M NaCl, 7%
SDS and 25mg/ml sonicated salmon sperm DNA and washed at 65°C in
0.1X SSC containing 0.1% sodium dodecyl sulfate and exposed to Kodak
BioMax MS film. Lambda phage clones hybridizing to the probe were
- 20 plaque purified by successive plating and rescreening. Ten positive clones
were purified and subjected to further analysis by hybridization to a 45 nt
oligonucleotide probe directed to the most 5' sequences known from the
original cDNA clone. This oligonucleotide was the reverse complement of
nucleotides 157-201 (Figure 2) and has the sequence 5'-CCAGGCGGCC
- 25 GCCATCTTGG AGACCGACAC TTTCTCGCCA CTTCC. Lambda phage
DNA was isolated by standard molecular biology techniques and
subjected to direct sequencing using fluorescent dideoxy cycle
sequencing on an ABI 373 sequencer.

- RACE: First strand DNA synthesis was performed using the rTth
- 30 thermal-stable polymerase system (Perkin Elmer). The following reagents
were combined in a 1.5 mL tube to give a 10 microliter volume: 1X

reverse transcription buffer, 1 mM MnCl_2 , 1.6 mM dNTP mix, 2.5U rTth polymerase, 100ng human hippocampus poly A⁺ RNA (Clontech), 10mM oligonucleotide (nt 429-452, Figure 2; 5'-GTTATGTTGG GTGCTGGAAA ACAG). The reaction was incubated at 70°C for 15 minutes and

5 immediately placed on ice. The Marathon cDNA synthesis kit (Clontech) was used for second strand cDNA generation. The entire 10 μ l from the first strand reaction was combined with the following reagents: 1X second strand buffer, 0.8 mM dNTP mix, 4X second strand cocktail (E.coli DNA polymerase I, E.coli DNA ligase, E.coli RNaseH), and dH₂O up

10 to a volume of 80 μ l. The tube was incubated at 16°C for 1.5 hours after which time T4 DNA polymerase (10U) was added and incubated for an additional 45 minutes at 16°C. To terminate the reaction, 4 μ l of 20X EDTA/glycogen (0.2M EDTA/2mg/ml glycogen) was added to the reaction mixes followed by a phenol/chloroform/isoamyl alcohol extraction to

15 remove enzymes and other impurities. The DNA was precipitated by adding 0.1X volume 3M Na acetate pH 5.2 and 2.5X volume reagent grade EtOH and place at -70°C. The DNA was washed once with 70% EtOH, dried down and resuspended in 10 μ l dH₂O. Half of the DNA was used for Marathon adaptor ligation to be used in subsequent RACE PCR

20 reactions following the Clontech protocol as follows: 5 μ l cDNA was added to 2 μ l (10mM) Marathon (5'-CTAATACGAC TCACTATAGG GCTCGAGCGG CCGCCCGGGC AGGT), 1X DNA ligation buffer and 1 μ l (1U) T4 DNA ligase. The reaction mix was incubated overnight at 16°C. The mix was diluted 1:50 for initial RACE reaction and combined in a

25 0.2mL PCR tube with the following: 40 μ l dH₂O, 1 μ l 10X Klentaq DNA polymerase (Clontech), 1 μ l (10mM) AP1 primer (5'-CCATCCTAAT ACGACTCACT ATAGGGC), 1 μ l (10mM) BBP1-specific primer (corresponding to nts. 187-209, Figure 2; 5'-CCAGACGGCCA GGCGGCCGCC AT), 5 μ l 10X Klentaq polymerase buffer, 1 μ l 10mM dNTP

30 mix, 1 μ l of diluted cDNA from above reaction. The following cycling conditions were performed using a Perking Elmer GeneAmp PCR system

2400 thermocycler: Denaturing cycle 94—C for 1 minute followed by 5 cycles of 30" at 94°C, 3' at 72°C, 5 cycles of 30" at 94°C, 3' at 70°C, followed by 25 cycles of 30" at 94°C, 3' at 68°C, with a final extension 7' at 72°C. This was followed by a nested RACE PCR reaction as follows: 40µl dH₂O, 1µl (1U) 10X AmplitaqGold DNA polymerase (Perkin Elmer), 1µl (10mM) AP2 primer (5'-ACTCACTATA GGGCTCGAGC GGC), 1µl (10mM) BBP1-specific primer (corresponding to nts. 172-194, Figure 2; 5'-GCCGCCATCT TGGAGACCGA CAC), 5µl 10X Amplitaq polymerase buffer, 1µl 10mM dNTP mix, 1µl of primary RACE product. The PCR cycling conditions were an initial denaturing cycle of 9' at 94°C, 25 cycles of 30" at 94°C, 30" at 68°C, 2' at 72°C, followed by a 72°C extension for 7'. The PCR product was run on a 1% agarose gel in 1XTBE buffer. The resulting 350 base pairs product was gel purified and directly cloned using the TA Cloning Kit (Invitrogen). Ligation mixes were transformed into OneShot Cells (Invitrogen) and plated on LB-ampicillin (100µg/ml) agar plates containing X-gal. Mini-prep DNA was obtained and examined by fluorescent dideoxy cycle sequencing on an ABI 373 sequencer.

Northern analyses. Human multiple tissue and multiple brain tissue mRNA Northern blots were obtained from Clontech (Palo Alto, CA). BBP1 sequences extending from the original fusion junction to the poly-A region were isolated on an EcoRI fragment from a TA clone derived from pEK196. β-actin DNA was provided by the manufacturer. Radiolabelled probes were produced from these DNAs using a random priming method to incorporate ³²P-dCTP (Pharmacia Biotech, Piscataway, NJ). Hybridizations were performed per manufacturer's (Clontech) instructions in Express Hyb Solution at 68°C. Blots were washed in 2x SSC (1X SSC is 0.15 M sodium chloride, 0.015 M sodium citrate), 0.05% SDS at room temperature, followed by two washes in 0.1 x SSC, 0.1% SDS at 50°C. Hybridization signals were visualized by exposure to Kodak BioMax film.

In situ hybridization. DNA templates for riboprobe synthesis were prepared by PCR using a plasmid clone containing the full length human BBP cDNA. A single riboprobe targeted to the 3' UTR of the cDNA was used. The probe sequences were checked versus the GenBank database to ensure that they only recognize the appropriate targets among all deposited sequences. To generate riboprobes for BBP1, a pair of oligonucleotide primers was designed to amplify a 275 base pairs region from the 3' UTR of the BBP1 cDNA and, in addition, add the promoter sequences for T7 (sense) and T3 (antisense) polymerase. These primers contained the following sequences: 5'-TAATACGACT CACTATAGGG TTAGAAGAAA CAGATTTGAG (forward); 5'-ATTAACCCTC ACTAAAGGGA CAAGTGGCAA CTTGCCTTTG (reverse). PCR products were gel purified on 1.5% low-melt agarose gels, and bands containing the products were excised, phenol and phenol-chloroform extracted, and ethanol precipitated. Pellet were dried and resuspended in 1X TE buffer (10 mM Tris-HCl, 1mM EDTA, pH 7.4). The APP riboprobe template consisted of a Ddel-XhoI fragment from the protein coding region, as described by Jacobsen et al. (1991). Fifty ng of DNA template was used for transcription reactions using (³⁵S)-CTP (New England Nuclear, Boston, MA) and the Riboprobe Gemini™ System (Promega, Madison, WI).

In situ hybridization histochemistry using sections of postmortem human hippocampus were performed as described previously (Rhodes, 1996). Sections were cut at 10 µm on a Hacker-Brights cryostat and thaw-mounted onto chilled (-20°C) slides coated with Vectabond reagent (Vector Labs, Burlingame, CA). All solutions were prepared in dH₂O treated with 0.1% (v/v) diethylpyrocarbonate and autoclaved. Sections were fixed by immersion in 4% paraformaldehyde in PBS (pH 7.4) then immersed sequentially in 2xSSC, dH₂O, and 0.1M triethanolamine, pH 8.0. The sections were then acetylated by immersion in 0.1M triethanolamine containing 0.25% (v/v) acetic anhydride, washed in 0.2xSSC, dehydrated in 50, 70 and 90% ethanol, and rapidly dried. One

ml of prehybridization solution containing 0.9M NaCl, 1mM EDTA, 5x Denhardt's, 0.25 mg/ml single-stranded herring sperm DNA (GIBCO/BRL, Gaithersburg, MD), 50% deionized formamide (EM Sciences, Gibbstown, NJ) in 10mM Tris, (pH 7.6), was pipetted onto each slide, and the slides

5 incubated for 3 hrs. at 50°C in a humidified box. The sections were then dehydrated by immersion in 50, 70, and 90% ethanol and air dried. Labeled riboprobes were added at a final concentration of 50,000 cpm/ μ l to hybridization solution containing 0.9M NaCl, 1mM EDTA, 1x Denhardt's, 0.1 mg/ml yeast tRNA, 0.1 mg/ml single-stranded salmon

10 sperm DNA, dextran sulfate (10%), 0.08% BSA, 10mM DTT (Boehringer Mannheim, Indianapolis, IN), and 50% deionized formamide in 10mM Tris (pH 7.6). The probes were then denatured at 95°C (1 min), placed on ice (5 min), and pipetted onto the sections and allowed to hybridize overnight at 55°C in a humidified chamber. The sections were subsequently

15 washed 1 x 45 min at 37°C in 2xSSC containing 10mM DTT, followed by 1 x 30 min at 37°C in 1xSSC containing 50% formamide, and 1 x 30 min at 37°C in 2xSSC. Single stranded and non-specifically hybridized riboprobe was digested by immersion in 10mM Tris pH 8.0 containing bovine pancreas RNase A (Boehringer Mannheim; 40 mg/ml), 0.5M NaCl,

20 and 1mM EDTA. The sections were washed in 2XSSC for 1 hr at 60°C, followed by 0.1XSSC containing 0.5% (w/v) sodium thiosulfate for 2 hrs. at 60°C. The sections were then dehydrated in 50, 70, 90% ethanol containing 0.3M ammonium acetate, and dried. The slides were loaded into X-ray cassettes and opposed to Hyperfilm b-Max (Amersham) for 14-

25 30 days. Once a satisfactory exposure was obtained, the slides were coated with nuclear-track emulsion (NTB-2; Kodak) and exposed for 7-21 days at 4°C. The emulsion autoradiograms were developed and fixed according to the manufacturer's instructions, and the underlying tissue sections were stained with hematoxylin. To assess nonspecific labeling, a

30 control probe was generated from a template provided in the Riboprobe Gemini™ System kit (Promega). This vector was linearized using Scal and

transcribed using T3 polymerase. The resulting transcription reaction generates two products, a 250 base and a 1,525 base riboprobe, containing only vector sequence. This control probe mixture was labeled as described above and added to the hybridization solution at a final concentration of 50,000 cpm/ μ l. No specific hybridization was observed in control sections, i.e., these sections gave a very weak uniform hybridization signal that did not follow neuroanatomical landmarks (data not shown).

Example 1: Cloning and Isolation BAP-binding protein (BBP1).

10 A yeast 2-hybrid (Y2H) genetic screen was developed to identify proteins which interact with human BAP₄₂, a 42 amino acid proteolytic fragment of APP which is considered to potentially be the more toxic aggregated form of BAP. BAP₄₂ was expressed fused to the yeast Gal4 DNA-binding domain and was also expressed as free peptide (Figure 1).

15 This strain was transformed with a human fetal brain cDNA Y2H library. A single clone, designated clone14 defined above, from approximately 10⁶ independent transformants, produced consistent reporter gene activation and contained a substantial open reading frame continuous with that of the GAL4 domain. The cDNA insert comprised 984 base pairs,

20 terminating in a poly-A tract. This sequence encoded 201 amino acids (SEQ ID NO: 2; amino acid residues 68 to 269) with two regions of sufficient length and hydrophobicity to transverse a cellular membrane.

 The library-derived plasmid was isolated from clone 14 and used to reconstruct Y2H assay strains. Examination of these strains

25 demonstrated that the BAP fusion protein specifically interacted with the clone 14 protein, although the response was weak. Since protein domains of strong hydrophobicity, such as transmembrane regions, inhibit Y2H responses (Ozenberger, unpublished data), clone 14 insert was truncated (hereinafter BBP1 Δ tm) to remove the region of strongest

30 hydrophobicity and retested for interactions with BAP. A much more robust Y2H response was observed with BBP1 Δ tm (Figure 2), supporting

the notion that the deleted sequences encode a potential transmembrane ("tm") anchor. The nucleotide sequence of Clone 14 was searched against GenBank; the BAP binding protein (BBP1) thus identified appeared to be novel.

5

Example 2: Isolation and confirmation of the 5' terminus of BBP1.

The BBP1 cDNA sequences contained in clone 14 described in Example 1, above, lacked the 5' end of the protein coding region as no potential initiating methionine codon was present. Multiple attempts at
10 conventional 5' RACE (rapid amplification of cDNA ends) utilizing a standard reverse-transcriptase only resulted in the addition of 27 nucleotides. These sequences included an ATG, but no upstream stop codon in the same translational reading frame to provide confidence that this was the initiating codon. A genomic cloning approach was initiated
15 to isolate the 5' terminus of the BBP1 gene.

Hybridization of a human genomic lambda library with a randomly-primed probe corresponding to 400 base pairs (bps) of the 5' sequence of clone 14 resulted in identification of 10 positive clones. Further characterization of these clones using a 45-base oligonucleotide probe
20 directed to the most upstream BBP1 sequence of clone 14 (and corresponding to the 5' upstream sequence of the 400 base pairs probe revealed that 6 of the 10 clones included the terminal 5' sequences contained within those previously identified. It was determined that the other 4 lambda clones represented other exons which were contained
25 within the original 400 base pairs randomly-primed cDNA-derived probe (data not shown). Direct cycle sequencing of lambda phage DNA from representative clones corresponding to the 5' end of BBP1 revealed ^a500 nucleotides upstream and overlapping with the sequence known for clone 14. This additional sequence potentially encodes 62 additional amino
30 acids upstream of the previously characterized MET before arriving at a MET preceded by an in-frame stop codon. Although there exist two MET

residues downstream from the furthest upstream MET, by standard convention we have tentatively defined the sequence of the amino terminus of the human BBP1 gene to include the first 5' MET which follows an in-frame stop codon. The entire coding region and deduced
5 protein sequence is shown in SEQ ID NOS:1 and 2. A plasmid (denoted BBP1-fl) containing this amino acid sequence has been deposited in the American Type Culture Collection having accession number 98617).

Since the 5' coding sequences were derived from a genomic library, there existed the possibility that this region contained introns.
10 This potentiality was investigated by two methods. First, a forward primer directed to the region of the 5' MET and a reverse primer within the original clone 14 were utilized to amplify sequences from brain cDNA as well as from genomic DNA. Products of identical size were generated from both samples, indicating the absence of introns within this region
15 and confirming the linkage of the upstream sequence with the original sequence. Secondly, cDNA sequences were isolated in modified 5' RACE experiments (see Materials and Methods, above) that were identical to those obtained from the genomic clone. These findings confirmed the upstream sequences (both from genomic and cDNA sources) and the lack
20 of introns in this region.

Example 3: Characterization of BBP1.

BBP1 sequences were compared to Genbank using the basic local alignment search tool (BLAST; Altschul et al., 1990). Two *Caenorhabditis*
25 *elegans* and one *Drosophila melanogaster* genomic sequence and a large number of human, mouse and other mammalian expressed sequence tags were identified. However, no complete cDNA sequences were available nor were any functional data attributed to the gene. The BBP1 protein and translations of available expressed sequence tags were aligned,
30 searched for conserved segments, and evaluated by the MoST (Tatusov et al., 1994) protein motif search algorithm. These analyses revealed a

potential evolutionary relationship to the G protein-coupled receptor family. Specifically, these analyses indicated that BBP1 contains two potential transmembrane (tm) domains equivalent to tm domains 3 and 4 of G protein-coupled receptors. The intervening hydrophilic loop contains
5 a well-characterized three amino acid motif, aspartate (D) or glutamate followed by arginine (R) and an aromatic residue (Y or F) (commonly referred to as the DRY sequence), that is conserved in almost all members of this receptor family and has been shown to serve as a molecular trigger for G protein activation (Acharya and Karnik, 1996). These data indicate
10 that BBP1 represents a novel protein potentially containing a functional module shared with members of the G protein-coupled receptor superfamily. Specifically, it appears that BBP1 retains the critical DRF sequence between two predicted tm domains, so may have the potential to couple to a G protein regulated signaling pathway.

15 Structural analysis of BBP1 indicated it contained a structural motif known to be a G α protein activating sequence in the related G protein-coupled receptors. Y2H assays demonstrating the interaction of BBP1 with various members of the G protein coupled receptors are illustrated in Figure 3. Based on structural predictions, BBP1 is depicted as
20 transversing a membrane twice with both termini in the luminal compartment. Other orientations cannot be entirely ruled out. The potential protein interactions described above were investigated in Y2H assays. Two overlapping portions of the BBP1 sequences contained in the BBP1 Δ tm clone were amplified and cloned into the Y2H vector pACT2
25 (expression plasmids pEK216 and pEK219, Table 2 and corresponding proteins BBP1 Δ N and BBP1 Δ C, Figure 4). The Δ C construct is lacking both tm domains; the Δ N construct encodes the first tm domain plus the preceding 52 amino acids. These fusion proteins were assayed with the BAP fusion protein and responses compared to those of strains expressing
30 the larger BBP1 Δ tm protein. These results suggest that a major determinant for the association with BAP is contained within the BBP1

region predicted to be topographically similar to BAP in the wild-type APP protein.

Example 4: Tissue distribution of human BBP1 expression.

5 Expression of BBP1 mRNA was evaluated as an initial step in elucidating the activity of the gene and its product. A major transcript of 1.25 kb was observed in all tissues (Figure 5A). There was a high level of expression in heart. Whole brain exhibited an intermediate level of expression. Samples derived from separate brain regions all exhibited
10 BBP1 expression (Figure 5B). Interestingly, limbic regions contained relatively greater amounts of BBP1 mRNA. These are the regions of the brain where BAP aggregation and associated neurotoxicity initially occur. Analysis of in situ hybridization autoradiograms obtained using a BBP1-specific riboprobe indicated that in human hippocampus and entorhinal
15 cortex, BBP1 mRNA is expressed in medium to large cells in a pattern consistent with expression in neurons as opposed to glial cells (Figure 6). Moreover, BBP1 mRNA is expressed in virtually all hippocampal and entorhinal neurons, i.e., there do not appear to be any real or laminar differences in the intensity of the hybridization signal. Interestingly, the
20 pattern of BBP1 expression was strikingly similar to the pattern observed using a riboprobe directed against mRNA for the amyloid precursor protein APP (Figure. 6). In summary, BBP1 mRNA was observed in all tissues and all brain regions examined. In situ analysis of BBP1 mRNA expression also revealed extensive expression in the hippocampus region.

25

Example 5: Cell line distribution of BBP1 expression.

 BBP1 expression was also investigated in numerous cell lines and data were extracted from dbEST, the collection of expressed sequence tags from the National Center for Biotechnology Information. Reverse-
30 transcription polymerase chain reaction (RT-PCR) methods were utilized to qualitatively assess BBP1 mRNA expression in cell lines commonly utilized

for recombinant protein expression as well as a variety of cancer cell lines. BBP1 was observed in hamster CHO and human HEK293 cells. Signals were observed in the embryonic stem cell line Ntera-2 and neuroblastoma lines IMR32 and SK-N-SH. BBP1 expression was observed

5 in cancer cell lines representing the following tissue origins: colon (Cx-1, Colo205, MIP101, SW948, CaCo, SW620, LS174T), ovarian (A2780S, A2780DDP), breast (MCF-7, SKBr-3, T47-D, B7474), lung (Lx-1, A5439), melanoma (Lox, Skmel30), leukemia (HL60, CEM), prostate (LNCAP, Du145, PC-3). A Northern blot probing mRNA isolated from the following

10 cancer cell lines demonstrated BBP1 expression in all samples: promyelocytic leukemia (HL-60), carcinoma (HeLa S3), chronic myelogenous leukemia (K-562), lymphoblastic leukemia (MOLT-4), Burkitt's lymphoma (Raji), colorectal adenocarcinoma (SW480), lung carcinoma (A549), and melanoma (G361).

15

Example 6: Selective interaction of BBP1 with human
BAP versus rodent BAP

There are three amino acid substitutions (G5R, F10Y and R13H) in the rodent BAP sequence compared to the human sequence. The rodent

20 peptide demonstrated reduced neurotoxicity and an absence of binding to human brain homogenates (Maggio et al., 1992). It was of interest, therefore, to evaluate the association of rodent BAP with BBP1 in the Y2H system. The sequence of human BAP in pEK162 was changed to encode the rodent peptide by oligonucleotide directed mutagenesis by

25 PCR, described above. The resultant plasmid, pEK240, was identical to the human BAP fusion protein expression plasmid utilized throughout the present invention except for the three codons producing the amino acid substitutions for the rodent peptide sequence. Interactions between BBP1 fusion protein and rodent and human BAP fusion proteins were compared

30 by Y2H bioassay. Strains expressing BBP1 and the rodent BAP failed to produce a growth response (Figure 7). This finding supports the premise

that BBP1 may serve as a specific mediator of the neurotoxic effects of BAP, and provides a mechanism to explain the reduced neurotoxicity of the rodent BAP. Importantly, these data also serve to illustrate the high degree of specificity of the BBP1/BAP interaction in the Y2H assays since the substitution of three amino acids was sufficient to completely abrogate the association (Figure 7).

Example 7: In vitro binding of radiolabeled beta-amyloid protein to BBP1 protein

Initially, the novel gene product, BBP1, expressed from a fetal brain library as a fusion protein, was shown to interact with beta-amyloid protein (BAP), also expressed as a fusion protein via a yeast 2 hybrid system. To confirm these initial findings, the potential binding of beta-amyloid protein to full-length BBP1 protein was investigated in an in vitro radioligand binding assay. Specifically, radiolabeled human beta-amyloid protein (1-42) was shown to bind with in vitro synthesized myc-tagged BBP1 protein, as evidenced by the ability to co-precipitate beta-amyloid protein with tagged-BBP1 protein. The details of the radioligand binding assay are described below.

Protein A agarose bead + secondary antibody complexes were generated by incubating 2.5 μ L ImmunoPurePlus immobilized Protein A (Pierce, Rockford, IL) with 10 mg AffiniPure rabbit a-mouse IgG (H + L) (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) in 50mL cold low salt binding buffer (50mM Tris pH7.6, 150mM NaCl, 2 mM EDTA 1%IGEPAL, and protease inhibitors [5 μ g/mL leupeptin, 5 μ g/mL aprotinin, 2 μ g/mL pepstatin A, 0.25mMPMSF]) with rotation overnight at 4°C. The beads were washed 4X with 1mL binding buffer and were resuspended in 1.25mL binding buffer to give a 50% slurry. In some experiments, a 250mL aliquot of this slurry was incubated in Superblock (Pierce) with rotation overnight at 4°C. The beads were washed 4X with 1mL Superblock and resuspended in 125 μ L Superblock.

The DNA template for in vitro transcription/translation of the BBP1 protein, including a Kozak consensus sequence and sequences encoding a myc epitope, EQKLISEEDL, directly upstream of the first methionine of BBP1 coding region, was inserted into the BamHI/EcoRI sites of

5 pSP64polyA vector (Promega, Madison, WI). The DNA template was, in part, PCR generated, utilizing the forward primer, 5' GCAGGATCCCCACCATGGAGCAGAAGCTGATCAGCGAGGAGGACCTGC ATATTTTAAAAGGGTCTCCCAATGTGA 3' and reverse primer, 5' TCACGGCCTCCGGAGCAGACGG 3' and PFU polymerase, according to

10 the manufacturer's conditions (Stratagene, La Jolla, CA). The PCR cycling conditions were an initial denaturing step at 95°C for 3min, 30 cycles of denaturation at 94°C for 30 sec, annealing at 65°C for 30 sec, elongation at 72°C for 1min 30sec, and followed by a final elongation at 72°C for 5 min. The amplicon was digested with BamHI + NotI and

15 ligated to the 3' end of BBP1, housed on a NotI/EcoRI fragment, which had been previously gel purified from the recombinant expression cassette.

Approximately 2.5μCi of disaggregated [Krishnamurthy, K. et al. (1998). Characterization of fibrillogenesis of amyloid peptide. Abstracts of the American Chemical Society, vol. , p. , 215th National Meeting and

20 Exposition, March 29-April 2, 1998, Dallas, Tx.] human [¹²⁵I]-Tyr-Ab₍₁₋₄₂₎ (American Radiolabeled Chemicals, Inc., St. Louis, MO) was incubated with 5-10mL of N-terminal c-myc tagged human BBP1 (1/5-1/10 reaction volume obtained using the TNT SP6 Coupled Reticulocyte Lysate System

25 [Promega, Madison, WI]) with rotation for ~6hrs at 4°C in a final volume of 1mL cold low salt binding buffer (see above). Two micrograms of mouse α-myc and 25mL of the Agarose protein A/rabbit α-mouse IgG complex (see above) were added to the reaction tube and incubated at 4°C overnight with rotation. Immune complexes were washed 4X with 1mL

30 binding buffer and resuspended in 25mL 2X Tricine loading dye (Novex, San Diego, CA) containing 5% β-Mercaptoethanol. Samples were boiled

for 5 minutes and immediately placed on ice for 15 minutes. The tubes were briefly spun at 2500 x g and the supernatant loaded on a 16% Tricine polyacrylamide gel (Novex, San Diego, CA) which was run at 50mA for ~90 min. The gel was soaked for 15 min in a drying solution
 5 composed of 20% acetic acid/10% methanol and dried at 80°C for 1 hr under vacuum. The dried gel was subjected overnight to a phosphorimager screen which was scanned for analysis with the Storm phosphorimager (Molecular Dynamics, Sunnyvale, CA).

Initial experiments attempting to co-immunoprecipitate radiolabeled
 10 BAP with myc-tagged BBP1 resulted in nonspecific binding of BAP when agarose protein A/secondary antibody complexes were prepared in low salt binding buffer, even in samples lacking BBP1 (see Fig. 8, lane 7). To reduce these non-specific interactions, the agarose protein A/rabbit a-mouse IgG was incubated/washed in blocking reagent prior to binding, as
 15 outlined above. This blocking procedure reduced non-specific Ab binding to near zero when all immunoprecipitation components were available except myc-tagged BBP1 (Fig. 8, lane 5). Radiolabeled human BAP₍₁₋₄₂₎ was able to complex with in vitro transcribed/translated myc-tagged human BBP1 after immunoprecipitating myc-tagged BBP1 with anti-myc
 20 antibody (Fig. 8, lane 9), as seen by a band consistent in size with Ab (Fig. 8, lane 3). These data are consistent with human BAP binding to myc-tagged human BBP1 in vitro and support the initial observation that BAP interacts with BBP1 in a yeast two-hybrid system.

25 **Example 8: Expression of recombinant BBP1 sensitizes
Ntera2 stem cells to β -amyloid peptide.**

A cultured cell system was utilized to investigate the effects of BBP1 expression on cellular sensitivity to BAP toxicity. Human Ntera-2 (Nt2) stem cells can be induced to differentiate into neuron-like cells (P.
 30 Andrews, Dev. Biol. 103:285-293, 1984). In that state, the cells exhibit a vulnerability to BAP that is similar in degree to that observed in primary neurons. Neurons affected by BAP exhibit characteristics of apoptosis

before dying (see C. Cotman and A. Anderson, *Mol. Neurobiol.* 10:19-45, 1995). An early indicator of apoptosis, namely, condensation of chromatin, was used as an indicator for cellular responses to BAP. The undifferentiated stem cells did not exhibit significant sensitivity under the experimental conditions used in these studies. However, Nt2 stem cells transfected with a BBP1 expression plasmid became markedly sensitive to applied BAP, supporting the premise that BBP1 may act as a mediator of the toxic effects of β -amyloid peptide. The details of the experiment are below.

BBP cDNAs were modified by polymerase chain reaction (PCR) for expression from the vector pcDNA3.1 (Invitrogen Corp., Carlsbad, CA). BBP1 cDNA was amplified from pBBP1-fl, adding a 5' EcoRI and a 3' Sall site for cloning. The PCR primers were 5' - TGGTGAATTC GAAAGTGTCTG GTCTCCAAG ATG G (+ strand) and 5' - CTTCGTCGAC TTA TGG ATA TAA TTG CGT TTT TC (- strand). The PCR product was digested with EcoRI + Sall and cloned into pcDNA3.1/EcoRI-XhoI to create pOZ363. Mutation of the arginine codon within the 'DRF' motif of the BBP1 cDNA was performed using the QuickChange system (Stratagene Co., La Jolla, CA). Oligonucleotides were synthesized and purified by Genosys Biotechnologies, Inc. (The Woodlands, TX). The R138 codon of BBP1 in pOZ363 was changed to a glutamate codon using the oligonucleotide 5' - GG TTG GGA GCA GAT GAA TTT TAC CTT GGA TAC CC and its exact reverse complement.

Human Ntera2 (Nt2) stem cells were maintained in Dulbecco's Modified Eagle's medium (high glucose) supplemented with 10% fetal bovine serum. Retinoic acid was utilized to differentiate cells to a neuronal phenotype as described by P. Andrews (*Dev. Biol.* 103:285-293, 1984). Expression constructs were introduced into stem cells by electroporation. The cells were split 1:2 the day before electroporation to ensure exponential growth for maximal survival and efficiency. On the day of electroporation the cells were treated with trypsin and washed two

times in phosphate buffered saline (PBS). They were resuspended at
1.3x10⁷ cells per 0.3 ml in RPMI 1640 with 10 mM dextrose and 0.1 mM
dithiothriitol. DNA amounts were 7.5 mg subject DNA with 2.5 mg
pEGFP-N1 (CLONTECH Laboratories, Palo Alto, CA) to monitor
5 transfection. Cells were pre-incubated for 10 mins on ice with DNA,
pulsed, and post-incubated for 10 min on ice. A GenePulser instrument
(BioRad Corp., Hercules, CA) was utilized with a cuvette gap of 0.4 cm,
voltage of 0.24 kV, and capacitance of 960 mF. Cells were plated in
standard 24-well plates. Approximately 24 hrs after transfection, growth
10 medium was replaced with medium containing the indicated concentration
of BAP. After incubation for 44 to 48 hrs, the chromatin-specific dye
Hoechst 33342 (Molecular Probes, Inc., Eugene, OR) was added to a
concentration of 10 ng/ml. Medium was removed after 10 min and cells
were washed with PBS. Cells were then fixed by immersion in PBS
15 containing 4% paraformaldehyde.

Forty-residue β -amyloid peptide was obtained from AnaSpec, Inc.,
San Jose, CA. Peptide was dissolved and stored in hexafluoro-
isopropanol at 1 mg/ml. Peptide was lyophilized by pervasion with
nitrogen, then resuspended in 1.155 ml cell growth medium and divided
20 into 0.13 ml aliquots in a 96-well plate. The plate was shaken at 500
rpm for 4 hrs. Samples were then combined and normalized to a final
BAP concentration of 50 mM. The same preparation of aggregated (or
aged) BAP utilized in the described experiments was also shown to be
toxic to primary hippocampal neurons (data not shown). Forty-two
25 residue β -amyloid peptide was obtained from Bachem Bioscience Inc. It
was dissolved directly in cell growth medium and added to experimental
samples. This preparation had no discernible effect on differentiated Nt2
neurons.

Cells were visualized on a Zeiss Axiovert fluorescent microscope
30 fitted with dichroic filters as follows. Hoechst dye visualization utilized
excitation at 330 microns, emission at 450; EGFP visualization with

excitation at 475, emission at 535. A minimum of 60 transfected (EGFP +) cells were scored per sample.

5 β -amyloid peptide exhibited substantial neurotoxicity in culture only after aging to produce fibrillar aggregates. Peptide freshly dissolved in media showed reduced potency. To investigate potential BBP1 effects on BAP-mediated toxicity, Nt2 stem cells were transfected with pEGFP or with pEGFP plus the BBP1 expression plasmid pOZ363 as described. These samples were treated with aged (toxic) or fresh (nontoxic) BAP as described above. Nt2 neurons, which exhibited a much greater sensitivity
10 to BAP than the undifferentiated stem cells, were treated at the same time to assess toxicity of the BAP preparations. After treatment for 48 hours, the chromatin of cells was stained with Hoechst dye to reveal nuclear structure and the cells were then fixed. The neurons treated with aged BAP exhibited a marked increase in condensed nuclei compared to those treated with the fresh BAP (Fig. 9), confirming the respective
15 toxicity of the two preparations. In contrast to the observations with Nt2 neurons, the stem cell controls exhibited only a small (not statistically significant) response to the toxic BAP (pEGFP; Fig. 9). However, transfection of the cells with the BBP1 expression plasmid resulted in a
20 significant ($P < 0.001$, Yates modified chi-squared test) increase in the frequency of condensed nuclei (Fig. 9). Importantly, the nontoxic preparation of BAP had no effect in the assay (Fig. 9), establishing that the BBP1-mediated response in these assays is specific for formulations of BAP that are also toxic to Nt2 neurons, and suggesting a correlation
25 between BBP1 and the mechanism of BAP toxicity. In a similar experiment, the toxic BAP was added to transfected Nt2 stem cells at varying concentrations (0, 0.5, 2, or 8 micromolar). Again, cells transfected with pEGFP alone exhibited no significant response to BAP treatment (Fig. 10). In contrast, cells transfected with a BBP1 expression
30 plasmid demonstrated a substantial, significant ($P < 0.005$), and dose-dependent sensitivity to the toxic BAP (Fig. 10).

It was predicted that the BBP1 protein might modulate the activity of heterotrimeric G proteins based on its structural relationship to known G protein-coupled receptors and the demonstration that its predicted cytosolic loop can associate with Ga proteins in a yeast 2-hybrid assay (Example 3, Fig. 3). It has been shown that substitution of the conserved arginine residue in the 'DRF' motif of 7-transmembrane domain G protein-coupled receptors attenuates their activity (E. Burstein, T. Spalding, and M. Brann, J. Biol. Chem. 273:24322-24327, 1998; W. Rosenthal et al., J. Biol. Chem. 268:13030-13033, 1993), and that substitution to glutamate (E) can completely eliminate agonist-mediated activation of G protein (P. Jones, C. Curtis, and E. Hulme, Eur. J. Pharmacol. 288:251-257, 1995). The BBP1 arginine-138 codon was mutated to a codon for glutamate and this BBP1 variant was examined for induction of sensitivity to BAP. This single amino acid substitution abrogated the effects of BBP1 (Fig.10), suggesting that the BAP sensitization affected by BBP1 in this system may be mediated through a G protein pathway.

It is clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and therefore are within the scope of the appended claims.

References

- Acharya, S., and Karnik, S. (1996). Modulation of GDP release from transducin by the conserved Glu134-Arg135 sequence in rhodopsin. *J Biol Chem* 271, 25406-25411.
- 5 Altschul, S., Gish, W., Miller, W., Myers, E., and Lipman, D. (1990). Basic local alignment search tool. *J Mol Biol* 215, 403-410.
- Chartier-Harlin, M., Crawford, F., Houlden, H., Warren, A., Hughes, D.,
10 Fidani, L., Goate, A., Rossor, M., Roques, P., Hardy, J., and Mullan, M. (1991). Early-onset Alzheimer's disease caused by mutations at codon 717 of the β -amyloid precursor protein gene. *Nature* 353, 844-846.
- DuYan, S., Chen, X., Fu, J., Chen, M., Zhu, H., Roher, A., Slattery, T.,
15 Zhao, L., Nagashima, M., Morser, J., Migheli, A., Nawroth, P., Stern, D., and Schmidt, A. (1996). RAGE and amyloid- β peptide neurotoxicity in Alzheimer's disease. *Nature* 382, 685-691.
- El Khoury, J., Hickman, S., Thomas, C., Cao, L., Silverstein, S., and
20 Loike, J. (1996). Scavenger receptor-mediated adhesion of microglia to β -amyloid fibrils. *Nature* 382, 716-719.
- Goate, A., Chartier-Harlin, M., Mullan, M., Brown, J., Crawford, F.,
Fidani, L., Giuffra, L., Haynes, A., Irving, N., James, L., Mant, R.,
25 Newton, P., Rooke, K., Roques, P., Talbot, C., Pericak-Vance, M., Roses, A., Williamson, R., Rossor, M., Owen, M., and Hardy, J. (1991). Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349, 704-706.
- 30 Hendriks, L., van Duijn, C., Cras, P., Cruts, M., van Hul, W., van Harskamp, F., Martin, J.-J., Hofman, A., and van Broeckhoven, C.

(1992). Presenile dementia and cerebral haemorrhage linked to a mutation at codon 692 of the β -amyloid precursor protein gene. *Nat Genet* 1, 218-221.

- 5 Jacobsen, J., Spruyt, M., Brown, A., Sahasrabudhe, S., Blume, A., Vitek, M., Muenkel, H., and Sonnenberg-Reines, J. (1994). The release of Alzheimer's disease β amyloid peptide is reduced by phorbol treatment. *J Biol Chem* 269, 8376-8382.
- 10 Jacobsen, J, Muenkel, H, Blume, A, and Vitek, M (1991). A novel species-specific RNA related to alternatively spliced amyloid precursor protein mRNAs. *Neurobiol of Aging* 12, 575-583.
- Kang, Y.-S., Kane, J., Kurjan, J., Stadel, J., and Tipper, D. (1990). Effects of expression of mammalian Ga and hybrid mammalian-yeast Ga proteins on the yeast pheromone response signal transduction pathway. *Mol Cell Biol* 10, 2582-2590.
- 15 Kang, Y.-S., Kane, J., Kurjan, J., Stadel, J., and Tipper, D. (1990). Effects of expression of mammalian Ga and hybrid mammalian-yeast Ga proteins on the yeast pheromone response signal transduction pathway. *Mol Cell Biol* 10, 2582-2590.
- LaFerla, F., Tinkle, B., Bieberich, C., Haudenschield, C., and Jay, G. (1995). The Alzheimer's A β peptide induces neurodegeneration and apoptotic cell death in transgenic mice. *Nature Genetics* 9, 21-30.
- 20 LaFerla, F., Tinkle, B., Bieberich, C., Haudenschield, C., and Jay, G. (1995). The Alzheimer's A β peptide induces neurodegeneration and apoptotic cell death in transgenic mice. *Nature Genetics* 9, 21-30.
- Lassmann, H., Bancher, C., Breitschopf, H., Wegiel, J., Bobinski, M., Jellinger, K., and Wisniewski, H. (1995). Cell death in Alzheimer's disease evaluated by DNA fragmentation in situ. *Acta Neuropathol* 89, 35-41.
- 25 Lassmann, H., Bancher, C., Breitschopf, H., Wegiel, J., Bobinski, M., Jellinger, K., and Wisniewski, H. (1995). Cell death in Alzheimer's disease evaluated by DNA fragmentation in situ. *Acta Neuropathol* 89, 35-41.
- Levy, E., Carman, M., Fernandez-Madrid, I., Power, M., Lieberburg, I., vanDuinen, S., Bots, G., Luyendijk, W., and Frangione, B. (1990). Mutation of the Alzheimer's disease amyloid gene in hereditary cerebral hemorrhage, Dutch type. *Science* 248, 1124-1126.
- 30 Levy, E., Carman, M., Fernandez-Madrid, I., Power, M., Lieberburg, I., vanDuinen, S., Bots, G., Luyendijk, W., and Frangione, B. (1990). Mutation of the Alzheimer's disease amyloid gene in hereditary cerebral hemorrhage, Dutch type. *Science* 248, 1124-1126.

- Loo, D., Copani, A., Pike, C., Whittemore, E., Walencewicz, A., and Cotman, C. (1993). Apoptosis is induced by β -amyloid in cultured central nervous system neurons. *Proc Natl Acad Sci USA* 90, 7951-7955.
- 5 Maggio, J., Stimson, E., Ghilardi, J., Allen, C., Dahl, C., Whitcomb, D., Vigna, S., Vinters, H., Labenski, M., and Mantyh, P. (1992). Reversible in vitro growth of Alzheimer disease β -amyloid plaques by deposition of labeled amyloid peptide. *Proc Natl Acad Sci USA* 89, 5462-5466.
- 10 Mullan, M., Crawford, F., Axelman, K., Houlden, H., Lilius, L., Winblad, W., and Lannfelt, L. (1992). A pathogenic mutation for probable Alzheimer's disease in the N-terminus of β -amyloid. *Nat Genet* 1, 345-347.
- 15 Murrell, J., Farlow, M., Ghetti, B., and Benson, M. (1991). A mutation in the amyloid precursor protein associated with hereditary Alzheimer disease. *Science* 254, 97-99.
- Nishimoto, I., Okamoto, T., Matsuura, Y., Takahashi, S., Okamoto, T.,
20 Murayama, Y., and Ogata, E. (1993). Alzheimer amyloid protein precursor complexes with brain GTP-binding protein Go. *Nature* 362, 75-79.
- Ozenberger, B., and Young, K. (1995). Functional interaction of ligands and receptors of the hematopoietic superfamily in yeast. *Mol Endocrinol*
25 9, 1321-1329.
- Rhodes K., Monaghan M., Barrezueta N., Nawoschik S., Bekele-Arcuri Z., Matos M., Nakahira K., Schechter L., and Trimmer J. (1996). Voltage-gated K⁺ channel beta subunits: expression and distribution of Kv beta 1
30 and Kv beta 2 in adult rat brain. *J Neurosci* 16, 4846-4860.

- Scheuner, D., Eckman, C., Jensen, M., Song, X., Citron, M., Suzuki, N., Bird, T., Hardy, J., Hutton, M., Kukull, W., Larson, E., Levy-Lahad, E., Viitanen, M., Peskind, E., Poorkaj, P., Schellenberg, G., Tanzi, R., Wasco, W., Lannfelt, L., Selkoe, D., and Younkin, S. (1996). A β 42(43) is
 5 increased in vivo by the PS1/2 and APP mutations linked to familial Alzheimer's disease. *Nat Med* 2, 864-870.

- Selkoe, D. (1997). Alzheimer's Disease: Genotypes, phenotype, and treatments. *Science* 275, 630-631.

10

- Smale, G., Nichols, N., Brady, D., Finch, C., and Jr, W. H. (1995). Evidence for apoptotic cell death in Alzheimer's disease. *Exp Neurol* 133, 225-230.

- 15 Tanzi, R., Gusella, J., Watkins, P., Bruns, G., George-Hyslop, P. S., vanKeuren, M., Patterson, D., Pagan, S., Kurnit, D., and Neve, R. (1987). Amyloid β protein gene: cDNA, mRNA distribution and genetic linkage near the Alzheimer locus. *Science* 235, 880-884.

- 20 Tatusov, R., Altschul, S., and Koonin, E. (1994). Detection of conserved segments in proteins : Iterative scanning of sequence databases with alignment blocks. *Proc Natl Acad Sci USA* 91, 12091-12095.

- Wade Harper, J., Adami, G. R., Wei, N., Keyomarsi, K., and Elledge, S. J.
 25 (1993). The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* 75, 805-816.

- Watt, J., Pike, C., Walencewicz-Wasserman, A., and Cotman, C. (1994). Ultrastructural analysis of β -amyloid-induced apoptosis in cultured
 30 hippocampal neurons. *Brain Res* 661, 147-156.

- Yamatsuji, T., Matsui, T., Okamoto, T., Komatsuzaki, K., Takeda, S.,
Fukumoto, H., Iwatsubo, T., Suzuki, N., Asami-Odaka, A., Ireland, S.,
Kinane, T., Giambarella, U., and Nishimoto, I. (1996). G protein-mediated
neuronal DNA fragmentation induced by familial Alzheimer's Disease-
5 binding mutants of APP. *Science* 272, 1349-1352.
- Yan, S., Fu, J., Soto, C., Chen, X., Zhu, H., Al-Mohanna, F., Collison, K.,
Zhu, A., Stern, E., Saido, T., Tohyama, M., Ogawa, S., Roher, A., and
Stern, D. (1997). An intracellular protein that binds amyloid- β peptide and
10 mediates neurotoxicity in Alzheimer's disease. *Nature* 389, 689-695.

11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624